

TEREBRA LINEOLATA, Sow. Tank. Cat. p. 76.

Buccinum voisin du *B. vittatum*.

DESH 1859 p 318

LINEOLATA

Sow 1825

EXCL

EXCLUENDAE

III. Anfractibus sulco postico nullo.

\*\* Labio interiore incrassato, elevato; testâ brevi.

OBS. Nassæ quodammodo affines; sed neque labium internum dilatatum, nec externum incrassatum.

Ter. lineolata, Sow. Wood, Suppl., f. 22.

GRA 34 p 63

Terebra lineolata G. B. Sowerby, 1825, Cat. shells coll. Tankerville, App. p. xxiii. This is a synonym of Bullia tranquebarica (Röding, 1798), family Nassariidae. 37-87  
EXC

18. Buccin linéolé. *Buccinum lineolatum*.

p 269

*B. testâ ovato-conicâ, læviusculâ, albido-cærulescente, lineolis fusco-nigris creberrimis interruptis seriâtim cinctâ; anfractibus convexis, supernè depressis; labro margine acuto, intus striato.*

?? BUCCINUM

Encyclop. pl. 400. f. 8. a. b.

Habite.... Mon cabinet. Longueur, 15 lignes.

LAM 1822

EXCLUDENDÆ.

T. LINEOLATA, Sowerby, l. c. p. 76.

HMS 446 p 176

1983. TEREBRA LINEOLATA.  
Chemn. iv. t. 155, f. 1463.  
T. testâ oblongâ, turritâ, lævi, albâ, lineolis longitudinalibus, subundatis, flavidis pictâ; anfractibus 7 rotundatis, basi sulcatis, margine superiore sulco valido prope suturam: long.  $1\frac{2}{10}$  lat.  $\frac{5}{10}$  unc.  
Obs. The figure of Chemn. above quoted is referred to by Dillwyn as a representation of *Buccinum vittatum*, which our shell resembles in general form; the sutures, however, are not crenulated, and it has only one groove, close to the upper edge of each volution. From Tranquebar. Sow 1825 App p xxiii

Excludendæ.

137. TEREBRA LINEOLATA, Sowerby, Tank. Cat.

*Buccinum lineolatum*, Wood, Index, Suppl.

*B. Bellangeri*, Kiener. HMS 1844 p 167

Lineolata (Terebra), Sow. Tank. Cat. App., p. 23.

= Bullia Bellangeri, Kr., Manual, iv,

39

TR40N1885