

EXCLENDAE

TEREBRA OBESA. Ter. testâ obeso-subulatâ, lævigatâ, albidâ, maculis fuscis longitudinalibus pallidè ornatâ; anfractibus paucis, subrotundatis, integris, ultimo biserialim maculato; spirâ obsoletè plicatâ; aperturâ oblongâ; columellâ truncatâ. Axis 6 lin.  
Hab. — ?  
Cab. Cuming. Unique.  
In this singular little shell the last whorl occupies nearly one half of the entire length.

H 44 A p 158

Species 140. (Mus. Cuming.)

TEREBRA OBESA. Ter. testâ cylindræo-subulatâ, albâ, maculis aurantiis amplis sparsim pictâ, anfractibus primis costatis, costis cito evanidis; aperturâ longâ, columellâ appressè arcuatâ, brevi.

THE STOUT TEREBRA. Shell cylindrically subulate, white, painted with a few large orange whorls convex, the first ribbed, ribs soon disappearing; aperture oblong, columella appressly a short.

HINDS, Pro. Zool. Soc. 1843, p. 158.

Hab. — ?

Resembling the preceding species in general form character, but painted with large blotches of pale-orange.

RV 1860



140.

HECALFII

94. T. OBESA. (pl. xlv. f. 106.) Hinds, l. c. p. 158.

Testâ obeso-subulatâ, lævigatâ, albidâ, maculis fuscis longitudinalibus pallidè ornatâ; anfractibus paucis subrotundatis, integris, ultimo biserialim maculato; spirâ obsoletè plicatâ; aperturâ oblongâ; columellâ truncatâ.

Polished and white, with a single series of pale spots; whorls entire, the inferior obese and smooth, the superior longitudinally plicate. Aperture oblong; columella slightly callous, truncate. No canal. I am only conversant with a single specimen, in the collection of Mr. Cuming, of this small but very distinct species.

Native country unknown.

H. 446 T. c. p 182 t xlv f 106



106.

POULSEN ?? VOIE A HASTATA DALL 1889 p 62

38. TEREBRA OBESA, Hinds.

Terebra obesa, Hinds, Thes. Conch. p. 182. no. 94. pl. 45. f. 106.

Hab. — ?

DESA 1859 p 283

146. obesa, Terebra - HINDS, 1844, P. Z. S. L., p. 158.

Hab. ? Coll. Cuming; unique! Size: 6 lin. (= 13.5 mm). Holotype no. 1968250: 13.6 mm. The labrum of the aperture is lirate in the type.

C69

T. OBESA, Hinds. Pl. 11, fig. 6.

Cylindrically subulate, whorls convex, the first ribbed, ribs soon disappearing; shining white, with longitudinal orange blotches, interrupted by a white band on the periphery.

Length, 12 mill.

Hab. unknown.



T. obesa Hinds

TRAYON 85 p 36

6. Terebra obesa, Hinds. Reeve, Conch. Icon., f. 140.

Obesa (Terebra), Hinds. Proc. Zool. Soc., 1843, p. 158, 116. 36

T 440 N 1885

Terebra obesa Hinds, 1844, Proc. Zool. Soc. Lond., pt. 10: 158. The outer lip of the aperture of the holotype is lirate, showing it does not belong in the Terebridae. 32 77

EXCL