

EXCUCIENDAE

PHALLUS  
G.M. 1791  
NOT A TEREBRA

Bosc 1801

**Phallus.** 146. *B. spirae* anfractibus longitudinaliter costatis basi futura rugosa, labro prominulo superne emarginato.  
*Martini. Conch. 4. t. 155. f. 1464. 1465.*  
*Schroet. Flusconch. p. 401. n. 194.*  
*Habitat in Indiae, an fluviis? testa tenui sub epidermide fusca flavicante, subfusca aut rufescente diaphana, costis albis, intus laevis; spirae anfractibus 9, costis subincurvis.*  
G.M. 1791 p. 3503

**BUCCINUM PHALLUS**, Gmel. p. 3503. no. 146.  
M. Pfeiffer dans son Index de Martini et Chemnitz rapporte au *Pleurotoma buccinoides* de Lamarek la figure de Martini dont Gmelin s'est servi pour le *B. phallus*.  
Bosc 1801 p. 319

There still remain to notice the following species, which were referred by the old authors to that section of *Buccinum* which is now regarded as synonymous with *Terebra*, but which there is much difficulty and doubt in referring to any shells at the present time in our collections:—*Buccinum murinum*, Linn.; *B. acus*, Chemnitz; *B. heciticum*, Gmelin; *B. bifasciatum*, Petiver; *B. concinnum*, Gualtier; *B. succinctum*, Chemnitz; *B. geminum*, Linn.; *B. proximatum*, Linn.; *B. monile*, Linn.; *B. sinuatum*, Linn.; and *B. phallus*, Chemnitz, is since known as *Pleurotoma buccinoides*.  
HDNS 4426108

**Vis phallus, Terebra phallus.**  
Les tours de spire avec des côtes longitudinales; la base de la suture inégale; la lèvre prominente et supérieurement émarginée.  
*Martini, Conch. tab. 155. fig. 1464 et 1465.*  
Se trouve dans l'Inde. Bosc 1801 p. 18

*Terebra phallus* Bosc, 1801, Hist. Nat. Coq. 5: 17. This is a synonym of *Clionella sinuata* (Born, 1778), family Turridae.  
Bosc 87