



x 13/10

H.A. 1859

97. *ACUS (EURYTA) TRILINEATA.

Euryta trilineata, A. Adams & Angas, P. Z. S. 1863, p. 418, pl. 37. f. 13.

An elegant fusiform shell, with the whorls longitudinally plicate and encircled with two or three brown thread-like lines. Dredged in deep water, near Port Jackson Heads. Length 7 lines.

ADAMS 1863 p 202

EXCLUENDAE

TRILINEATA

ADAMS & ANGAS 1863

1863

EXC

817—*trilineata* Adams & Angas, *Euryta*, Proc. Zool.

Soc., 1863, p. 418, pl. 37, f. 13. H.A. 1859

Trilineata (*Euryta*), Ad. and Ang. P. Z. S., 1863, p. 418, pl. xxxvii, f. 13, 38



T. 24 trilineata Adams & Ang.

T. TRILINEATA, Adams and Angas. Pl. 12, fig. 24.

Thin, slender, with long, narrow aperture, distantly plicate and encircled with thread-like lines; yellowish white, with longitudinal pale undulating chestnut markings, a series of chestnut spots near the suture, and on the body-whorl three revolving chestnut lines, one of which appears also on the spire.

Length, 14 mill.

TR 40 N 85 p 38

New South Wales, Australia.

24. *Terebra trilineata*, Ad. and Angas. Zool. Proc., t. 37, f. 13, 1863.

Three-lined Auger (*Parviterebra trilineata*). Thin, slender, distantly plicated on surface, encircled with thin, thread-like lines. Aperture long, narrow. Yellow-white, with longitudinal, undulating pale chestnut markings and chestnut spots near the sutures and on body whorl. About 1/2 inch long, inhabits New South Wales, Victoria, Tasmania and South Australia. Probably also south-west Australia.

JACOB 1864

NOT in B. & C 87
I DO NOT KNOW IF IT HAS
EVER BEEN ASSIGNED
TO THE GENUS TEREBRA!

1. EURYTA TRILINEATA, Ad. & Ang. (Pl. XXXVII. fig. 13.)

E. testa turrito-fusiforimi, gracili; spira quam apertura longiore; albida, fasciis longitudinalibus undulatis fuscis, et serie macularum fuscicarum ad suturas ornata, lineisque tribus transversis filiformibus ad anfractum ultimum succincta; anfractibus 9, planiusculis, longitudinaliter plicatis, plicis in medio anfractuum subnodulosis, in anfractu ultimo obsolete, transversim sulcatis, sulcis subdistantibus; apertura angusta; labio antice violascente tincto.

Long. 7 lines, lat. 1 1/2 line.

Hab. Port Jackson (Coll. Angas.).

This species of *Euryta* (a group separated from *Terebra* on account of the spirally rolled axis and Mitriform aspect) differs from the others described in being more slender, and in the whorls being encircled with thread-like lines.

ANGAS 1864 p 418



13

13. *Euryta trilineata*



13

NOT THE ORIGINAL DESCRIPTION !!