

97. T. TUBEROSA. (pl. xlv. f. 99.) *Hinds*, l. c. p. 152.



Testâ turrîto-subulatâ, saturatè fulvâ, asperâ, nitidâ; anfractibus rotundatis, longitrorsum costatis, supernè cingulo numerosè tuberculato; costis brevibus, nodulosis striis decussantibus; columellâ contortâ.

In this characteristic species the girdle consists of a number of small tubercles, superior in number to the vertical ribs. H. 446. T. C. p. 183 t. xlv f. 99.

EXCULENDAE

TUBEROSA

HINDS 1844

1844

145. TEREBRA TUBEROSA, Hinds.

Terebra tuberosa, Hinds, *Thes. Conch.* p. 183. no. 97, pl. 45. f. 99.

Hab. — ?

A la juger par la figure, cette coquille semblerait un Cérîte dont l'ouverture aurait été mutilée ou serait restée imparfaite. D 254 1857 p 303

Tuberosa (*Terebra*), Hinds. *Proc. Zool. Soc.*, 1843, p. 152; *Thes. Conch.*, 183, t. 45, f. 99, 40

T. TUBEROSA, Hinds, *Hab.*? The figure appears to represent a *Cerithium* with imperfect aperture. T 2404 15 p 40

-T. (? *Strioterebrum*) *tuberosa* HINDS, 1844: *Proc. Zool. Soc. London* for 1843: 291; Fig.: HINDS in SOWERBY; *Thes. Conch.*, 1844, T., pl. 45, fig. 99.

This species has not been among the specimens examined. B 61466

Terebra tuberosa Hinds, 1844, *Proc. Zool. Soc. Lond.* pt. 11: 152. 1844, Hinds in Sowerby, *Thes. Conch.* 1: 183, pl. 45, fig. 99. The type is lost, and the figure suggests this is possibly a juvenile *Cerithium*. H. 446 T. C. p. 183 3277

TEREBRA TUBEROSA. *Ter. testâ turrîto-subulatâ, saturatè fulvâ, nitidâ; anfractibus rotundatis, longitrorsum costatis, supernè cingulo numerosè tuberculato; costis brevibus; nodulosis striis decussantibus; columellâ contortâ.* Axis 11 lin.

Hab. Ticao, Philippines.

Cab. Cuming. Unique.

In this characteristic species the girdle consists of a number of small tubercles, superior in number to the vertical ribs.

H. 446 p 152